

The Moore bound for Spectral Radius

Vladimir Nikiforov

Department of Mathematical Sciences, University of Memphis,
Memphis TN 38152, USA

February 2, 2008

Abstract

Let G be a graph with n vertices, m edges, girth g , and spectral radius μ . Then

$$\mu^2 \leq \begin{cases} m & \text{if } g = 4 \\ n - 1 & \text{if } g \geq 5. \end{cases}$$

Keywords: Moore bound, spectral radius, irregular graph, degree sequence

What is the maximum spectral radius of a graph of order n and girth g ? Equivalently, what is the minimum order of a graph with girth g and spectral radius μ . For d -regular graphs the spectral radius is equal to d ; thus, the Moore bound (see, e.g., [2], p.180 and [4] for a general survey) shows that the order of such graphs is at least

$$n(d, g) = \begin{cases} 1 + d + d(d-1) + \cdots + d(d-1)^{r-1} & \text{if } g = 2r + 1 \\ 2(1 + (d-1) + \cdots + (d-1)^{r-1}) & \text{if } g = 2r. \end{cases}$$

Alon, Hoory, and Linial [1] extended this result, showing that every graph with girth g and *average degree* d has at least $n(d, g)$ vertices.

The bound changes radically when the average degree is replaced by the spectral radius.

Theorem 1 *Let G be a graph with n vertices, m edges, girth g , and spectral radius μ . Then*

$$\mu^2 \leq \begin{cases} m & \text{if } g = 4 \\ n - 1 & \text{if } g \geq 5. \end{cases}$$

If $g = 4$, then $\mu^2 = m$ holds if and only if G is a complete bipartite graph possibly with isolated vertices. If $g \geq 5$ then $\mu^2 = n - 1$ holds if and only if G is either a Moore graph of diameter 2 or a star.

For $g = 4$ this result was obtained in [6] and extended in another direction in [5]. The case $g \geq 5$ was given in [3].

References

- [1] N. Alon, S. Hoory, N. Linial, The Moore bound for irregular graphs, *Graphs Combin.* **18** (2002), 53–57.
- [2] N. Biggs, *Algebraic graph theory*, Cambridge University Press, Cambridge, second edition, 1993.
- [3] O. Favaron, M. Mahéo, J.-F. Sacle, Some eigenvalue properties in graphs (conjectures of Graffiti. II), *Discrete Math.* **111** (1993), 197–220.
- [4] M. Miller, J. Širáň, Moore graphs and beyond: A survey of the degree/diameter problem, www.combinatorics.org/Surveys/ds14.pdf.
- [5] V. Nikiforov, Some inequalities for the largest eigenvalue of a graph, *Combin. Probab. Comput.* **11** (2002), 179–189.
- [6] E. Nosal, Eigenvalues of Graphs, Master’s thesis, University of Calgary, 1970.